


LETTER TO THE EDITOR

Comment on 'Perinatal mortality after the Fukushima accident'

To cite this article: Hagen Scherb *et al* 2019 *J. Radiol. Prot.* **39** 647

View the [article online](#) for updates and enhancements.



IOP | IPEM

Series in Physics and Engineering in Medicine and Biology

Your publishing choice in medical physics,
biomedical engineering and related subjects.

Start exploring the collection and download the first chapter
of every title for free.

Letter to the Editor



CrossMark

Comment on ‘Perinatal mortality after the Fukushima accident’

Sir,

In their Letter to the Editor [1], Körblein and Küchenhoff discussed in some detail our publication [2] on perinatal mortality in Japan before and after the Fukushima nuclear power plant accidents. Employing annual data from 2002 to 2015 and a slightly different stratification of the Japanese prefectures, Körblein and Küchenhoff essentially confirmed our findings of long-term perinatal mortality increases relative to the prevailing stable secular downward trends. Our analyses were originally based on monthly data from 2001 to 2014. By the end of the year 2018, further official monthly perinatal data of the Japanese prefectures for the years 2016 and 2017 became available. Thus, it is now possible to extend our initiating analyses by three more years of data, namely from January 2015 through December 2017, which amounts to a considerable enhancement of statistical power and precision. Figures 1 and 2 show the monthly perinatal mortality (2002–2017) in the six severely contaminated prefectures Fukushima, Gunma, Ibaraki, Iwate, Miyagi, and Tochigi as well as in the three moderately contaminated prefectures Chiba, Saitama, and Tokyo, respectively. The jumps in perinatal mortality in January 2012 presented in figures 1 and 2 corroborate, firstly, Körblein and Küchenhoff’s results and, secondly, our preliminary findings based on the initial three years after Fukushima. Importantly, also after the Chernobyl accident long-term relative increases in stillbirths and perinatal mortality were reported along with exposure-dependent increases in congenital malformations [3, 4]. Consistent with these observations after major nuclear accidents, Kaori Murase *et al* [5] demonstrate an increasing number of surgeries for severe congenital heart malformations throughout Japan after Fukushima. Murase *et al* suggest indirectly that one of the causes of perinatal death includes severe congenital heart diseases. We recommend and hope that responsible scientists and institutions will continue to document and scrutinize the prevalence of perinatal deaths and congenital malformations in Japan more comprehensively over longer periods of time and stratified by exposure.

Sincerely,

Hagen Scherb, Kuniyoshi Mori, Keiji Hayashi

Hagen Scherb^{1,4} , Kuniyoshi Mori² and Keiji Hayashi³¹ Helmholtz Zentrum München, German Research Center for Environmental Health, Germany² Higashiosaka Health Center 4-3-22 Iwatachou, Higashiosakacity, Japan³ Hayashi Children’s Clinic, Osaka, JapanE-mail: hagen.scherb@gmail.com⁴ Author to whom any correspondence should be addressed.

Perinatal mortality in Fukushima, Gunma, Ibaraki, Iwate, Miyagi, Tochigi; 4-months peak from March 2011 and jump in January 2012




Figure 1. Monthly perinatal mortality (2002–2017) in six severely contaminated prefectures Fukushima, Gunma, Ibaraki, Iwate, Miyagi, and Tochigi; logistic regression model (solid line) adjusted for direct earthquake and tsunami effects and allowing for a delayed jump (+10 months) after Fukushima; the broken line indicates the null-effects/null-hypotheses model; jump in January 2012 with jump odds ratio (OR) 1.188 (1.085, 1.301), $P = 0.0002$.

Perinatal mortality in Chiba, Saitama, Tokyo; 4-months peak from March 2011 and jump in January 2012




Figure 2. Monthly perinatal mortality (2002–2017) in three moderately contaminated prefectures Chiba, Saitama, and Tokyo; logistic regression model (solid line) adjusted for direct earthquake and tsunami effects and allowing for a delayed jump (+10 months) after Fukushima; the broken line indicates the null-effects/null-hypotheses model; jump in January 2012 with jump odds ratio (OR) 1.106 (1.035, 1.183), $P = 0.0031$.

ORCID iDs

Hagen Scherb  <https://orcid.org/0000-0002-2730-5619>

References

- [1] Körblein A and Küchenhoff H 2017 Perinatal mortality after the Fukushima accident *J. Radiol. Prot.* **37** 800
- [2] Scherb H, Mori K and Hayashi K 2016 Increases in perinatal mortality in prefectures contaminated by the Fukushima nuclear power plant accident in Japan *Medicine (Baltimore)* **95** e4958
- [3] Scherb H, Weigelt E and Brueske-Hohlfeld I 1999 European stillbirth proportions before and after the Chernobyl accident *Int J Epidemiol* **28** 932–40
- [4] Scherb H and Weigelt E 2003 Congenital malformation and stillbirth in Germany and Europe before and after the Chernobyl nuclear power plant accident *Environ. Sci. & Pollut. Res. Special Issue 1* 17–125
- [5] Murase K, Murase J and Mishima A 2019 Nationwide increase in complex congenital heart diseases after the Fukushima nuclear accident *J. Am. Heart Assoc.* **8** e009486