

CONFERENCE ON THE HUMAN RIGHT TO A SAFE AND HEALTHFUL ENVIRONMENT AND THE RESPONSIBILITIES UNDER INTERNATIONAL LAW OF OPERATORS OF NUCLEAR FACILITIES

Salzburg, October 20-23, 2005

Summary of Long-Term Risks Created by Prolonged Contact with Low-Level Radioactivity

Inge Schmitz-Feuerhake, German Society of Radiation Protection, Cologne, Germany
e-mail: ingesf@uni-bremen.de

Summary

The International Commission on Radiological Protection ICRP has developed risk estimates about health detriments in populations exposed by low doses of ionizing radiation. They were derived from findings in the Japanese A-bomb survivors and are specified to three kinds of effects: 1) hereditary diseases in the descendants, 2) Cancer mortality, 3) Developmental defects by in utero exposure. Numerous findings in contaminated regions show that the effects are underestimated by the ICRP risk factors by several orders of magnitude in cases of chronic low dose exposure to incorporated radioactivity. Alternatively, the risk estimates of the European Committee on Radiation Risk ECRR are presented.

The most serious radiation effects by radioactivity – hereditary defects in the descendants of exposed parents – had been already detected in the twenties of the last century by the later nobel prize winner Herman Joseph Muller. He concluded from his investigations in drosophila that also low dose exposures and thus also the natural background radiation are mutagenous. In the thirties already, the idea arose that cancer is initiated by a single cell transformation, a “somatic” mutation. Therefore Muller concluded that there is also no harmless dose range for cancer induction (1).

Later on, the International Commission on Radiological Protection ICRP developed the term of the “stochastic” radiation effect from this. If a great collective is exposed by a small dose, one cannot predict which individual person will suffer from a radiation damage, only a probability is derivable. The amount of diseases increases with the accumulated dose, but after halving the dose there remains still an elevated effect. Therefore, no “threshold” exists i.e. a dose range without risk.

A third group of radiation effects which has to be considered in the low dose region was also known already at the time of Muller. These are teratogenesis effects i.e. damages which are caused by exposure in utero. They may manifest as cancer in childhood or as malformations and other developmental distortions, also as fetal loss, stillbirths, and infant deaths. Kind and frequency of the damage for a given exposure depend on the stage of development. In the former days of radiation research the fetus was thought to be the most radiosensitive system.

The Chernobyl accident, at the latest, has shown that several other serious diseases occur after chronic low dose irradiation. Distortions of the central nervous system are observed, especially mental illnesses (2), as well as malfunctions of other organs. It is often supposed that distortions of the immune system are the cause of the observed diseases because deviations in the corresponding parameters were measurable. It was further confirmed that radiation-induced lens opacities (cataracts) have to be counted among the stochastic effects.

The geneticist Muller warned after the second world war of deteriorating the genetic pool of mankind by environmental radioactivity. He was therefore uninvited as speaker at the Atomic Conference of Geneva in 1955 where the large-scale so-called peaceful use of nuclear energy was initiated. Since then, those scientists were preferred and selected as experts by the authorities who declared the handling of man-made huge amounts of radioactivity to be practicable and safe. The anti-nuclear movement was initiated by scientists whose conclusion was that the ruling opinions about the effects of radioactivity were wrong and dangerous, as e.g. was expressed by John Gofman and Arthur Tamplin (3), former advisers of the U.S. Atomic Energy Commission AEC which had been established for the promotion of nuclear energy application.

The normative board for the evaluation of radiation risks and the proposal of dose limits is the ICRP. Although the commission derives only recommendations these are applied by all Western and Eastern industrial nations. It followed a committee which had been founded in 1928 by radiological societies of several countries for the purpose of developing standards for radiation protection in the medical field. Therefore, it is traditionally obliged to the interests of the users. Since 1950, in the period of the Cold War and the development of nuclear energy consumption, it grew up to great importance. Technically, its recruitment remained the task of the radiological societies, de facto it recruits itself (4).

Critics of the official radiation protection criteria have therefore always questioned the arguments and results of the ICRP which completely contradict many established findings, especially in regions contaminated by radioactivity. The reference population which is used by the ICRP to play down the effects are the Japanese A-bomb survivors of Hiroshima and Nagasaki. The research programs in these are dominated by American scientists.

The problems of such comparison are not only lying in several peculiarities which reduce the general validity of the Japanese data. It lies also in the dosimetry system which was created by the ICRP to make all exposure conditions comparable. The Japanese survivors were irradiated by extremely high-energetic penetrating gamma rays, the radioactive fallout is neglected in the dose estimation. The dose in Sv is defined as an absorbed energy per tissue mass (dimension Joule/kg) multiplied with a so-called weighting factor depending on the kind of radiation: gamma/beta or alpha radiation. For estimating the dose in cases of radionuclide incorporation it is necessary to know the amount of substance deposited in the specific tissue and the course of retention from there. The ICRP has developed dose factors in Sv per inhaled or ingested Bq of the radionuclide which are intended to meet all individual conditions. The uncertainties of these dose factors may amount to several orders of magnitude (5, 6). Nevertheless are they used – to the disadvantage of the affected persons – quite similarly as natural constants and prescribed in the official rules of radiation protection e.g. to decide whether the dose limits near nuclear installations are kept.

In order to overcome an endless discussion about the parameters used in the ICRP models Busby chose another method to derive realistic estimates. He was followed by an European Committee on Radiation Risk (ECRR) founded in 1998 which sees itself as an alternative of the ICRP (7). Its risk estimates are based on observed evidence in the low-dose region for those effects where the basic mechanisms of induction are known.

The risk estimates of the ICRP about health effects after low-dose exposure are shown in Table 1 (8, 9) and compared to the evaluation of the ECRR.

Table 1 Health effects by chronical low-dose irradiation of a population

	Hereditary diseases	Cancer mortality	Teratogenic effects (in utero exposure)	Morbidity except from tumours
ICRP Risk estimate	130 cases per 10 ⁴ Sv	500 deaths per 10 ⁴ Sv (5 % per Sv)	No effect below 100 mSv	No effect
Evaluation by the ECRR	Underestimated by factor 100-2000	Underestimated by factor 100-2000	Cancer Malformations Mental retardation Mental disorders Down 's syndrome Childhood morbidity Stillbirths Infant deaths Spontaneous abortions Low birth weight	Manifold

The value 10⁴Sv in columns 2 and 3 expresses a unity of the collective dose which is the sum of all individual doses in a collective. The risk estimate by the ICRP in column 2 means, that 130 additional cases of hereditary diseases are expected if a population of 10.000 persons is exposed by 1 Sv or a population of 10 million people by 1 mSv. Such relation can also be considered as an individual risk. 500 deaths per 10⁴Sv corresponds to a probability of 5 % for suffering from cancer death after exposure by 1 Sv.

For dose comparison it may be useful to remember that the dose limit – after recommendation of the ICRP - for a population exposed by emissions from nuclear facilities is 1 mSv per year.

It is shown in the ECRR report that the ICRP either simply ignores numerous results in the scientific literature about low level effects or neglects them for doubtful reasons. A most grotesque attitude is shown by their evaluation of teratogenic effects. The mainstream science has accepted in the meantime – after resistance for decades – that the cancer risk is real for children who were prenatally exposed to diagnostic x-rays and thus to very low doses. This damage has to be attributed therefore to the stochastic effects without threshold dose. The ICRP also adopts this not without distaste, in contrast, however, to this formal acceptance they postulate their threshold dose of 100 mSv which is not even proved in the A-bomb survivors, their preferred reference collective.

Numerous studies after diagnostic x-raying and in regions contaminated by radioactivity – especially by the Chernobyl accident – have shown a variety of teratogenic effects which are listed in column 4 of table 1 (10).

Regarding genetic diseases not only the ICRP but also other international and national committees of radiation protection refer exclusively to the A-bomb survivors where no significant effects were observed in the first generation of descendants. The ICRP interpretes their risk estimate (Table 1, column 1) therefore as an estimate on the safe side which would probably not occur in reality. The many results about genetically induced cancer diseases which were initiated by the debate about the leukemias near the British reprocessing plant Sellafield are declared to be not plausible in view of the knowledge of Hiroshima and Nagasaki.

Table 2 is taken from the ECRR report (7) and contains informations about European nuclear facilities which are showing elevated cancer rates after 1983 when the Sellafield leukemias had been detected. In these contaminated regions one has to consider as well as for numerous

findings after Chernobyl that there will be a cumulative induction in both exposed parents as well as by prenatal and postnatal irradiation of the children.

Table 2 Studies establishing excess leukaemia and cancer risk in children living near nuclear sites, data from ECRR-Report 2003 (7)

Nuclear Site	Year	Notes	Reference
^a Sellafield/Windscale, UK	1983	Well studied by ^f COMARE: high level of discharge to atmosphere and sea	(11)
^a Dounreay, UK	1986	Well studied by ^f COMARE: high level of discharge to atmosphere and sea	(12)
^a La Hague, France	1993	Particle discharge to atmosphere and sea: ecological and case control studies	(13)
^c Aldermaston/Burghfield, UK	1987	Well studied by ^f COMARE: particle discharge to atmosphere and rivers	(14,15)
^b Hinkley Point, UK	1988	Discharges to offshore and bank	(16)
^d Harwell, UK	1997	Discharges to atmosphere and river	(17)
^e Birkenfeld, Germany	1990	Discharges to atmosphere and drinking water	(18, 19)
^{b,d} Geesthacht, Germany	1992	Discharges to atmosphere and Elbe river	(20-22)
^d Jülich, Germany	1996	Discharges to atmosphere and river	(23)
^b Barsebaeck, Sweden	1998	Discharges to atmosphere and sea	(24)

^aReprocessing plants discharging to sea; ^bNuclear power station discharge to sea or river; ^cAtomic weapon and nuclear material fabrication plants; ^dAtomic research with discharges to local rivers, ^eUranium waste, ^fCommittee on Medical Aspects of Radiation in the Environment, U.K.

Results about radiation-induced cancers in children which were only preconceptionally exposed are compiled in Table 3. The dose estimates were derived by the authors or reckoned by the writer. From these and the relative risks the doubling dose was derived (the dose which elevates the rate of diseases by the same amount as is given by the spontaneous rate).

This compilation shows that the interpretation of the Sellafield leukaemias by Martin Gardner (25) as genetically induced was not at all new and unexpected as was claimed by ICRP and other committees. Rather, such connection had been found by a variety of researchers before.

Studies in children of parents which were exposed by Chernobyl radioactivity have shown that not only cancer is genetically induced in the next generation by low level exposure but also malformations, metabolic diseases, mental disorders, and Down's syndrome (35).

Table 3 Radiation-induced hereditary effects/Cancer in childhood after preconceptional low-dose exposure

Exposed collective	Disease	Gonadal Dose mSv	Relative Risk	Doubling dose mSv
Seascale fathers 1990 (25)	Leukaemia + lymphoma “ “	200	7	29
all stages of spermatogenesis		10	7	1.4
6 months before conception			1.9	
Sellafield workers 2002 (26)			3.2	
Occupational exposure W.Cumbria 1991 (27)				
Preconceptional X-ray diagnostics				
Fathers 1966 (28)	Leukaemia	5*	1.3	3.8
Fathers 1988 (29)	Leukaemia	3-30	1.4-3.9	
Fathers 1994 (30)	Leukaemia		3.8	
Mothers 1958 (31)	Leukaemia	5*	1.7	2.9
Mothers 1966 (28)	Leukaemia	5*	1.7	2.9
Mothers 1973 (32)	Leukaemia	5*	1.4	3.6
Mothers 1980 (33)	Cancer	3*	2.6	1.2
Occupational exposure 1984 (34)	Cancer		2.7	

* Dose values reckoned by writer

1. Muller, H.J.: Über die Wirkung der Röntgenstrahlung auf die Erbmasse. *Strahlentherapie* 55, 1936, 207-224
2. Loganovsky, K.N., Loganovskaja, T.K.: Schizophrenia disorders in persons exposed to ionizing radiation as a result of the Chernobyl accident. *Schizophr. Bull.* 26, 2000, 751-753
3. Gofman, John W., Tamplin, Arthur R.: *Poisoned Power. The Case against Nuclear Power Plants.* Rodale Press inc., Emmaus, Pa. 18049, 1971
4. Morgan, K.Z.: ICRP risk estimates – an alternative view. In Jones, R.R., Southwood, R. (Eds.), *Radiation and Health. Biological Effects of Low Level Exposure to Ionizing Radiation.* John Wiley & Sons 1987, p. 125-154
5. Fairlie, I.: Uncertainties in doses and risks from internal radiation. *Medicine, conflict and survival* 21, 2005, 111-126
6. Schmitz-Feuerhake, I.: Bewertung neuer Dosisfaktoren. In Dannheim, B. et al.: *Strahlengefahr für Mensch und Umwelt. Bewertungen der Anpassung der deutschen Strahlenschutzverordnung an die Forderungen der EU-Richtlinie 96/29/Euratom.* Berichte des Otto Hug Strahleninstituts Nr. 21-22, 2000, p. 55-74
7. ECRR, European Committee of Radiation Risk: *Health Effects of Ionising Radiation Exposure at Low Doses for Radiation Protection Purposes. 2003 Recommendations of the ECRR.* Eds. Busby, C. et al., Brussels 2003
8. ICRP, Int. Commission on Radiological Protection: *1990 Recommendations of the International Commission on Radiological Protection.* ICRP-Publ.60, Ann. ICRP 21, No.1-3, 1991
9. ICRP, Int. Commission on Radiological Protection: *Biological effects after prenatal irradiation (embryo and fetus).* ICRP Publication 90. *Annals of the ICRP* 33, No.1-2, 2003
10. Schmitz-Feuerhake, I.: Wie verlässlich sind die Grenzwerte? Neue Erkenntnisse über die Wirkung inkorporierter Radioaktivität. *Strahlentelex* Nr. 442-443 v. 2.6.05, S. 1-6
11. Independent Advisory Group. *Investigation of the possible increased incidence of cancer in West Cumbria.* London: HMSO, 1984. Black report.
12. Committee on the Medical Aspects of Radiation in the Environment: *Second Report. Investigations of the possible increased incidence of leukaemia in young people near the Dounreay Nuclear Establishment, Caithness, Scotland.* London: HMSO, 1988
13. Pobel, D., Viel, J.-F.: Case-control study of leukaemia among young people near La Hague nuclear reactor reprocessing plant: the environmental hypothesis revisited. *Br. Med. J.* 314, 1997, 101-106
14. Bithell, J.F., Dutton, S.J., Draper, G.J., Neary, N.M.: Distribution of childhood leukaemias and non-Hodgkin's lymphomas near nuclear installations in England and Wales. *Br. Med. J.* 309, 1994, 501-505
15. Barton, C.: The incidence of childhood leukaemia in West Berkshire. *Med. Confl. Surv.* 17, 2001, 48-55

16. Ewings, P.D., Bowie, C., Phillips, M.J., Johnson, S.A.: Incidence of leukaemia in young people in the vicinity of Hinkley Point nuclear power station. *Br. Med. J.* 299, 1989, 289-293
17. Busby, C., Cato, M.S.: Death rates from leukaemia are higher than expected in areas around nuclear sites in Berkshire and Oxfordshire. *Br. Med. J.* 315, 1997, 309
18. Hoffmann, W.: Inzidenz maligner Erkrankungen bei Kindern und Jugendlichen in der Region Ellweiler, Rheinland-Pfalz. Epidemiologie und Biologische Dosimetrie zur Ermittlung möglicher Belastungspfade. Diss. University of Marburg, Germany. Shaker, Aachen 1993
19. Hoffmann, W., Kranefeld, A., Schmitz-Feuerhake, I.: Radium-226-Contaminated Drinking Water: Hypothesis on an exposure pathway in a population with elevated childhood leukemia. *Environm. Health Perspectives* 101 (1993) Suppl. 3, 113-115
20. Schmitz-Feuerhake, I., Dannheim, B., Heimers, A., Oberheitmann, B., Schröder, H., Ziggel, H.: Leukaemia in the proximity of a German boiling water reactor: evidence of population exposure by chromosome studies and environmental radioactivity. *Health Persp.* 105, Suppl.6, 1997, 1499-1504
21. Schmitz-Feuerhake, I., Mietelski, J.W., Gaca, P.: Transuranic isotopes and ⁹⁰Sr in attic dust in the vicinity of two nuclear establishments in northern Germany. *Health Physics* 84, 2003, 599-607
22. Schmitz-Feuerhake, I., Dieckmann, H., Hoffmann, W., Lengfelder, E., Pflugbeil, S., Stevenson, A.F.: The Elbmarsch leukemia cluster: are there conceptual limitations in controlling immission from nuclear establishments in Germany? *Arch. Environ. Contamination Toxicol.* 49, 2005, 589-601
23. Kuni, H.: A cluster of childhood leukaemia in the vicinity of the German research reactor Jülich. In Schmitz-Feuerhake, I., Schmidt, M. (Eds.), *Radiation Exposures by Nuclear Facilities*. Gesellschaft für Strahlenschutz e.V., Berlin 1998, p. 251-255
24. Andersen, H., Moeller, T.: *Cancerinsidens omkring Barsebaecks Kaernkraftwerk, 1997*, Lund, Regionala Tumoregister Universitetssjukhuset
25. Gardner, M.J., Snee, M.P., Hall, A.J., Powell, A.J., Downes, S., Terrell, J.D.: Results of case-control study of leukaemia and lymphoma among young people near Sellafield nuclear plant in West Cumbria. *Brit. Med. J.* 300, 1990, 423-429
26. Dickinson, H.O., Parker, I.: Leukaemia and non-Hodgkin's lymphoma in children of male Sellafield radiation workers. *Int. J. Cancer* 99, 2002, 437-444
27. McKinney, P.A., Alexander, F.E., Cartwright, R.A., Parker, L.: Parental occupations of children with leukemia in west Cumbria, north Humberside, and Gateshead. *Br. Med. J.* 302, 1991, 681-687
28. Graham, S., Levin, M.L., Lilienfeld, A.M. et al.: Preconception, intrauterine, and postnatal irradiation as related to leukemia. *Natl. Cancer Inst. Monogr.* 19, 1966, 347-371
29. Shu, X.O., Gao, Y.T., Brinton, L.A., Linet, M.S., Tu, J.T., Zheng, W., Fraumeni, J.F.: A population-based case-control study of childhood leukemia in Shanghai. *Cancer* 62, 1988, 635-644
30. Shu, X.O., Reaman, G.H., Lampkin, B., Sather, H.N., Pendergrass, T.W., Robison, L.L.: Association of paternal diagnostic x-ray exposure with risk of infant leukemia. *Cancer Epidemiol., Biomarkers & Prevention* 3, 1994, 645-653
31. Stewart, A., Webb, J., Hewitt, D.: A survey of childhood malignancies. *Br. Med. J.* i, 1958, 1495-1508
32. Natarajan, N., Bross, I.D.J.: Preconception radiation leukemia. *J. Med.* 4, 1973, 276-281
33. Shiono, P.H., Chung, C.S., Myriantopoulos, N.C.: Preconception radiation, intrauterine diagnostic radiation, and childhood neoplasia. *J. Natl. Cancer* 65, 1980, 681-686
34. Hicks, N., Zack, M., Caldwell, G.G., Fernbach, D.J., Falletta, J.M.: Childhood cancer and occupational radiation exposure in parents. *Cancer* 53, 1984, 1637-1643
35. Yablokov, A. et al: *Compilation of health effects after the Chernobyl accident*. ECRP publication, in preparation