

**RADIATION-INDUCED EFFECTS IN HUMANS AFTER IN UTERO EXPOSURE:
CONCLUSIONS FROM FINDINGS AFTER THE CHERNOBYL ACCIDENT**

INGE SCHMITZ-FEUERHAKE*

Department of Physics, University of Bremen, Germany

Abstract. A threshold dose of 100 mSv is assumed for radiation-induced teratogenic effects in publication No. 90 of the ICRP (2003) which is based on the findings in the Japanese A-bomb survivors. A variety of observations about congenital malformations, fetal loss, stillbirths, and infant deaths, as well as Down's syndrome after the Chernobyl release demand to point out the incompleteness of the Japanese data. Radiation-induced developmental effects by Chernobyl, especially in great distances, are usually denied because of the low values of the estimated human exposures. Biological dosimetry in contaminated regions, however, shows that physical dose estimations lead to considerable underestimations of the true exposure. The assumptions about teratogenic effects by incorporated radionuclides have to be revised.

Keywords: radiation-induced malformations, perinatal mortality; Chernobyl effects in children; dosimetry of incorporated radioactivity, biological dosimetry

Introduction

The evaluation of radiation risks by international radiation protection committees is based on the findings in the Japanese A-bomb survivors. The effects observed there after prenatal exposure were mental retardation and reduced head size while no other significant detriment was detected. The time between the 8th and 15th week of gestation is thought to be the period of risk.

As was pointed out by different researchers the Japanese data suffer, however, from several restrictions which limit their suitability as a general base for deriving radiation risks. One point is a probable and proven severe selection bias because of the catastrophic situation after the bombing. Another objection which must be stressed especially considering perinatal effects is the fact that the investigations of the Radiation Effects Research Foundation (RERF) in Hiroshima did not begin earlier than 5 years after the catastrophe when the RERF research institute was established there. The completeness of the data must therefore be doubted.

The reduced selection of assumed effects and the assertion of a threshold dose as high as 100 mSv by the ICRP contradicts former findings in experimental research (Table 1) and several observations in humans already before the Chernobyl accident. Tables 2-5 list the results of studies in regions affected by fallout from the latter.

TABLE 1. Minimum dose below 100 mSv which showed significant effects after in utero x-ray exposure in experimental studies (data taken from Fritz-Niggli, 1997)

	Dose mSv	Days after conception	Effects	Reference
Mice	10	8	Cumulated developmental defects	Michel, Fritz-Niggli 1977
	50	0.5	Death of the embryo	Rugh, Grupp 1959
	50	0.5; 1.5	Death of the embryo, polydactyly	Ohzu 1965
	50	7.5	Death of the embryo, skeletal malformations	Jacobsen 1966
Rats	10	18	Reflex distortions	UNSCEAR 1986
	50	0.4; 0.7	Fetal death	Roux et al. 1983

* Retired; Peter-Michels-Str. 54, 50827 Cologne, Germany; e-mail: ingesf@uni-bremen.de

TABLE 2. Observed increase of congenital malformations after in utero exposure by the Chernobyl accident

Country	Effects	References
Belarus National Genetic Monitoring Registry	Anencephaly, spina bifida, cleft lip and/or palate, polydactyly, limb reduction defects, esophageal atresia, anorectal atresia, multiple malformations	Lazjuk et al. 1997
Belarus Highly exposed region of Gomel	Congenital malformations	Bogdanovich 1997; Savchenko 1995
Chechersky district of the Gomel region	Congenital malformations	Kulakov et al. 1993
Mogilev region	Congenital malformations	Petrova et al. 1997
Brest region	Congenital malformations	Shidlovskii 1992
Ukraine Polesky district of the Kiev region	Congenital malformations	Kulakov et al. 1993
Lygyny region		Godlevsky, Nasvit 1998
Turkey	Anencephaly, spina bifida	Akar et al. 1988/89; Caglayan et al. 1990; Güvenc et al. 1993; Mocan et al. 1990
Bulgaria, region of Pleven	Malformations of heart and central nervous system, multiple malformations	Moumdjiev et al. 1992
Croatia	Malformations by autopsy of stillborns and cases of early death	Kruslin et al. 1998
Germany German Democratic Republic, Central registry	Cleft lip and/or palate	Zieglowski, Hemprich 1999
Bavaria	Cleft lip and/or palate Congenital malformations	Scherb, Weigelt 2004 Korblein 2003a, 2004; Scherb, Weigelt 2003
Annual Health Report of West Berlin 1987	Malformations in stillborns	Strahlentelex 1989
City of Jena (Registry of congenital malformations)	Isolated malformations	Lotz et al. 1996

TABLE 3. Observed increase of stillbirths, infant deaths, spontaneous abortions, and low birth weight after in utero exposure by the Chernobyl accident

Country	Effects	References
Belarus Selected regions	Perinatal deaths*)	Petrova et al. 1997
Chechersky District near Gomel	Perinatal deaths	Kulakov et al. 1993
Gomel region	Perinatal deaths	Korblein 2003a,b
Ukraine Polessky District near Kiev	Perinatal deaths, reduced birth rate**), premature births	Kulakov et al. 1993
Lygny region	Early neonatal deaths	Godlevsky, Nasvit 1998
Zhitomir oblast, Kiev region, Kiev City	Perinatal deaths, reduced birth rate	Korblein 2003a,b
Europe: Greece, Hungary, Poland, Sweden	Stillbirths	Scherb et al. 1999b, 2000b, 2003
Poland	Infant mortality	Korblein 2003a
Norway	Spontaneous abortions	Ulstein et al. 1990
Hungary	Low birth weight	Czeisel 1988
Finland	Premature births among malformed children Reduced birth rate Stillbirths	Harjulehto et al. 1989 Harjulehto et al. 1991 Scherb, Weigelt 2003
Germany Total (FRG + GDR)	Perinatal deaths	Korblein, Kuchenhoff 1997; Scherb et al. 2000a,2003
Southern Germany	Early neonatal deaths	Lüning et al. 1989
Bavaria	Perinatal deaths, stillbirths Reduced birth rate	Grosche et al. 1997; Scherb et al. 1999a, 2000a, 2003 Korblein 2003a

*) Perinatal deaths summarize stillbirths and deaths in the first 7 days from birth

**) Reduced birth rate is considered as a measure for spontaneous abortions

TABLE 4. Increase of Down's syndrome after in utero exposure by the Chernobyl accident

Region	Results	References
Belarus National Genetic Monitoring Registry	Excess 1987-1994 ca. 17 % Excess peak in January 1987	Lazjuk et al. 1997 Zatsepin et al. 2004
Western Europe	Beginning 1 y after the accident, reaching 22% within 3 years	Dolk et al. 1999
Sweden	„Slight“ excess in most exposed areas (30 %)	Ericson, Kallen 1994
Scotland, Lothian region (0.74 million inhabitants)	Excess peak in January 1987 (2-fold significant)	Ramsay et al. 1991
South Germany	Elevation found by investigations of amniotic fluid	Sperling et al. 1991
Berlin West	Excess peak in January 1987	Sperling et al. 1991, 1994

TABLE 5. Observed health defects in children after in utero exposure by the Chernobyl accident except malformations and Down's syndrome

Belarus Selected regions	Mental disorders Speech-language disorders, mental retardation	Kondrashenko et al. 1996 Kolominsky et al. 1999
Chechersky District near Gomel	Diseases of respiratory organs, blood, circulation etc.	Kulakov et al. 1993
Stolin District in Brest region	Diseases of respiratory organs, glands, blood, circulation, digestive organs	Sychik, Stozharov 1999a,b
Belarus, Ukraine, Russia	Mental retardation and other mental disorders	Kozlova et al. 1999
Ukraine Polesky District near Kiev	Diseases of respiratory organs, blood, circulation etc.	Kulakov et al. 1993
Rovno Province	Childhood morbidity	Ponomarenko et al. 1993
Immigrants to Israel from contaminated areas	Asthma	Kordysh et al. 1995

Estimated exposures from the Chernobyl accident

In spite of the overwhelming evidence of effects observed also in countries far away from the Chernobyl event they are ignored by international radiation protection committees. They assume that the exposures of the population are much too low to generate teratogenic effects. Indeed, the physical dose estimates resulted in mean effective life-time exposures in large regions of Europe and in Turkey below 1.2 mSv (UNSCEAR 1988). The highest average dose for a subregion in the first year after the accident is derived to 2 mSv in Belarus.

These estimates show, however, large discrepancies in comparison with biological dosimetry. Investigations of unstable and stable chromosome aberrations in the lymphocytes of persons in the contaminated regions have been done by a variety of research groups in rather large collectives

directly after the accident or some years later. Dicentric chromosomes can be considered as radiation-specific and most sensitive because of their very low and nearly constant rate in unexposed persons (Hoffmann and Schmitz-Feuerhake, 1999).

It is a general experience that the observed rates of dicentric chromosomes after Chernobyl are considerably higher – by 1 or 2 orders of magnitude – than would be expected from physically derived dose estimates. This evaluation is possible although the dose-effect relationships in cases of incorporated radioactivity are not known. For low dose homogenous exposure by low LET irradiation the rate of dicentrics can be considered as dose-proportional which follows from studies in the range of background exposure. In European countries far from the Chernobyl site the exposure of the tissues except the thyroid is assumed to be mainly generated by ^{137}Cs and ^{134}Cs which distribute homogeneously inside the body. The whole body doubling dose for dicentrics by homogenous low LET radiation is about 10 mSv (Hoffmann and Schmitz-Feuerhake, 1999). Elevations of dicentrics in persons which are higher than 2-fold would therefore mean that the whole body dose exceeds 10 mSv. Such elevations have been found manifold after Chernobyl. Findings for Austria, Germany and Norway are shown in Table 6.

A remarkable finding in many of the chromosome studies is that they report an overdispersion of the dicentrics and the occurrence of multiaberrant cells (Bochkov and Katosova 1994; Hille et al. 1995; Salomaa et al. 1997; Scheid et al. 1993, Sevan`kaev et al. 1993; Stephan and Oestreicher 1989; Verschaeve et al. 1993). This is an indication for a relevant contribution of incorporated α -activity which is not considered adequately in the physical dose estimates.

TABLE 6. Chromosome aberrations in lymphocytes of persons living in West European regions contaminated by Chernobyl releases; dics dicentric chromosomes, cring centric rings

Region	Sample	Date of study	Method	Results mean elevation	Reference	Remarks
Salzburg, Austria	17 adults	1987	dics+crings	circ 4-fold	Pohl-Rüling et al. 1991	
Germany, southern regions	29 children + adults	1987-1991	dics+crings	circ 2.6-fold	Stephan, Oestreicher 1993	physical dose estimate <0.5 mSv
Norway, selected regions	44 reindeer sames 12 sheep farmers	1991	dics+crings	10-fold	Brogger et al. 1996	physical dose estimate 5.5 mSv

Conclusions

The data in tables 2-6 show that neither the exposure of populations affected by the Chernobyl accident nor the effects have been evaluated correctly up to now. It must be discussed, however, that some of the authors interpret their somatic findings as genetically induced. In cases of continual exposure by radioactivity it is, indeed, not always to decide whether the damage occurred in utero or by preconceptional mutation. Intrauterine deaths and infant deaths as well as malformations are also inducible by irradiation of the germ-cells in fathers or mothers. Rugh (1962) has already stated that the appearance of certain malformations of the brain is quite similar after preconceptional and in utero irradiation.

The studies in the A-bomb survivors did not show significant genetic effects. Investigations in the offspring of a large cohort of Sellafield workers, however, which were initiated after the leukaemia findings in the proximity of this British nuclear reprocessing plant, confirm a rise of stillbirths and congenital anomalies caused by preconceptional exposure of the fathers (Parker et al. 1999). The cited Chernobyl studies may therefore include some overlap of transgenerational and teratogenic effects. Although this conclusion demands further scientific effort for differentiation the fetus must be

considered as most vulnerable by low dose exposures as was assumed in the early times of radiation research.

- Akar, N., Cavdar, A.O., and Arcasoy, A., 1988, High incidence of Neural Tube defects in Bursa, Turkey, *Paediatric and Perinatal Epidemiol.* **2**:89-92.
- Akar, N., Ata, Y., and Aytekin, A.F., 1989, Neural Tube defects and Chernobyl? *Paediatric and Perinatal Epidemiol.* **3**:102-103.
- Bochkov, N.P., and Katosova, L.D., 1994, Analysis of multiaberrant cells in lymphocytes of persons living in different ecological regions, *Mutat. Res.* **323**:7-10.
- Bogdanovich, I.P., 1997, Comparative analysis of the death rate of children, aged 0-5, in 1994 in radiocontaminated and conventionally clean areas of Belarus, in: *Medicobiological effects and the ways of overcoming the Chernobyl accident consequence*, Collected book of scientific papers dedicated to the 10th anniversary of the Chernobyl accident, Minsk-Vitebsk, p. 4.
- Brogger, A., Reitan, J.B., Strand, P., and Amundsen, I., 1996, Chromosome analysis of peripheral lymphocytes from persons exposed to radioactive fallout in Norway, *Mutat. Res.* **361**:73-79.
- Caglayan, S. Kayhan, B., Montesoglu, S., and Aksit, S., 1990, Changing incidence of neural tube defects in Aegean Turkey, *Paediatric and Perinatal Epidemiol.* **4**:264-268.
- Czeisel, A.E.; and Billege, B., 1988, Teratological evaluation of Hungarian pregnancy outcomes after the accident in the nuclear power station of Chernobyl, *Orvosi Hetilap* **129**:457-462 (Hungarian).
- Dolk, H., Nichols, R., and a EUROCAT Working Group, 1999, Evaluation of the impact of Chernobyl on the prevalence of congenital anomalies in 17 regions of Europe, *Int. J. Epidemiol.* **28**:941-948.
- Ericson, A., and Kallen, B., 1994, Pregnancy outcome in Sweden after Chernobyl, *Environ. Res.* **67**:149-159.
- Fritz-Niggli, H., 1997, *Strahlengefährdung/Strahlenschutz*, 4th ed., Hans Huber, Bern, Switzerland.
- Godlevsky, I., and Nasvit, O., 1998, Dynamics of health status of residents in the Lugyny district after the accident of the ChNPS, in: *Research activities about the radiological consequences of the Chernobyl NPS accident and social activities to assist the sufferers by the accident*, T. Imanaka, ed., Research Reactor Institute, Kyoto University, KURRI-KR-21, pp.149-156.
- Grosche, B., Irl, C., Schoetzau, A., and van Santen, E., 1997, Perinatal mortality in Bavaria, Germany, after the Chernobyl reactor accident, *Rad. Environ. Biophys.* **36**:129-136.
- Güvenc, H., Uslu, M.A., Güvenc, M., Ozkici, U., Kocabay, K., and Bektas, S., 1993, Changing trend of neural tube defects in Eastern Turkey, *J. Epidemiol. Community Health* **47**:40-41.
- Harjulehto, T., Aro, T., Rita, H., Rytomaa, T., and Saxen, L., 1989, The accident at Chernobyl and outcome of pregnancy in Finland, *Brit. Med. J.* **298**:995-997.
- Harjulehto, T., Rahola, T., Suomela, M., Arvela, H., and Saxén, L., 1991, Pregnancy outcome in Finland after the Chernobyl accident. *Biomed & Pharmacother* **45**:263-266.
- Hille, R., Wolf, U., Fender, H., Arndt, D., and Antipkin, J., 1995, Cytogenetic examination of children in Ukraine, In *Radiation Research 1895-1995. Proceed. 10th Int. Congr. Radiation Res. Würzburg*, U. Hagen et al., ed., Germany.
- Hoffmann, W., and Schmitz-Feuerhake, I., 1999, How radiation-specific is the dicentric assay? *J. Exp. Analysis Environ. Epidemiol.* **2**:113-133.
- Jacobson, L., 1966, Radiation-induced effects in mouse foetuses, *Acta radiol.* **254**:82
- Kolominsky, Y., Igumnov, S., and Drozdovitch, V., 1999, The psychological development of children from Belarus exposed in the prenatal period to radiation from the Chernobyl atomic power plant, *J. Child. Psychol. Psychiatry* **40**:299-305.
- Kondrashenko, V.G. et al., 1996, Mental disorders caused by Chernobyl, in: *Report on the 3rd Int. Congress "World after Chernobyl"*, Minsk, 1996, cited in Nesterenko, V.B., *Chernobyl Accident. Radiation Protection of Populatio*, Republic of Belarus Institute of Radiation Safety "Belrad", Minsk 1998.
- Korblein, A., and Kuchenhoff, H., 1997, Perinatal mortality in Germany following the Chernobyl accident, *Rad. Environ. Biophys.* **36**:3-7.
- Körblein, A., Säuglingssterblichkeit nach Tschernobyl, 2003a, *Berichte Otto Hug Strahleninstitut* **24**:6-34.
- Korblein, A., 2003b, Strontium fallout from Chernobyl and perinatal mortality in Ukraine and Belarus, *Radiats. Biol. Radioecol.* **43**:197-202.
- Körblein, A., 2004, Fehlbildungen in Bayern nach Tschernobyl, *Strahlentelex No.* **416-417**:4-6.

- Kordysh, E.A., Goldsmith, J.R., Quastel, M.R., Poljak, S., Merkin, L., Cohen, R., and Gorodischer, R., 1995, Health effects in a casual sample of immigrants to Israel from areas contaminated by the Chernobyl explosion, *Environ. Health Persp.* **103**:936-941.
- Kozlova, I.A., Niagu, A.I., and Korelev, V.D., 1999, The influence of radiation to the child mental development, *Zh. Nevrol. Psikiatr. Im. SS Korsakova* **99**:12-16 (Russ.).
- Kruslin, B., Jukic, S., Kos, M., Simic, G., and Cviko, A., 1998, Congenital anomalies of the central nervous system at autopsy in Croatia in the period before and after Chernobyl, *Acta Med. Croatica* **52**:103-107.
- Kulakov, V.I., Sokur, T.N., Volobuev, A.I., Tzibulskaya, I.S., Malisheva, V.A., Zikin, B.I., Ezova, L.C., Belyaeva, L.A., Bonartzev, P.D., Speranskaya, N.V., Tchesnokova, J.M., Matveeva, N.K., Kaliznuk, E.S., Miturova, L.B., and Orlova, N.S., 1993, Female reproduction function in areas affected by radiation after the Chernobyl power station accident, *Environ Health Persp.* **101, Suppl. 2**:117-123.
- Lazjuk, G.I., Nikolaev, D.L., and Novikova, I.V., 1997, Changes in registered congenital anomalies in the Republic of Belarus after the Chernobyl accident, *Stem Cells* **15, Suppl. 2**:255-260.
- Lotz, B., Haerting, J., and Schulze, E., 1996, Veränderungen im fetalen und kindlichen Sektionsgut im Raum Jena nach dem Reaktorunfall von Tschernobyl, Oral presentation at the International Conference of the Society for Medical Documentation, Statistics, and Epidemiology, Bonn, Germany (available on request).
- Lüning, G., Scheer, J., Schmidt, M., and Ziggel, H., 1989, Early infant mortality in West Germany before and after Chernobyl, *Lancet II*:1081-1083.
- Michel, C., Fritz-Niggli, H., 1977, Radiation damage in mouse embryos exposed to 1 rad X-rays or negative pions, *Fortschritte Röntgenstrahlen* **127**:276-280.
- Mocan, H., Bozkaya, H., Mocan, Z.M., Furtun, E.M., 1990, Changing incidence of anencephaly in the eastern Black Sea region of Turkey and Chernobyl, *Paediatric and Perinatal Epidemiol.* **4**:264-268.
- Moumdjiev, N., Nedkova, V., Christova, V., Kostova, Sv., 1992, Influence of the Chernobyl reactor accident on the child health in the region of Pleven, Bulgaria, in: *20th Int. Congr. Pediatrics Sept. 6-10, 1992 in Brasil*, p.57. Cited by Akar, N., Further notes on neural tube defects and Chernobyl (Letter), *Paediatric and Perinatal Epidemiol.* **8**, 1994, 456-457.
- Ohzu, E., 1965, Effects of low-dose X-irradiation on early mouse embryos, *Radiat. Res.* **26**:107.
- Parker, L., Pearce, M.S., Dickinson, H.O., Atkin, M., Craft, A.W., 1999, Stillbirths among offspring of male radiation workers at Sellafield nuclear processing plant, *Lancet* **354**: 1407-1414
- Petrova, A., Gnedko, T., Maistrova, I., Zafranskaya, M., Dainiak, N., 1997, Morbidity in a large cohort study of children born to mothers exposed to radiation from Chernobyl. *Stem Cells* **16, Suppl. 2**: 141-150
- Pohl-Rüling, J., Haas, O., Brogger, A., Obe, G., Lettner, H., Daschil, F., Atzmüller, C., Lloyd, D., Kubiak, R., and Natarajan, A.T., 1991, The effect on lymphocyte chromosomes of additional burden due to fallout in Salzburg (Austria) from the Chernobyl accident, *Mutat. Res.* **262**:209-217.
- Ponomarenko, V.M., Nagornaia, A.M., Proklina, T.L., Litvinova, L.A., Stepanenko, A.V., Sytenko, E.R., and Osnach, A.V., 1993, The morbidity in preschool children living on the territory of Rovno Province subjected to radioactive contamination as a result of the accident at the Chernobyl Atomic Electric Power Station, *Lik Sprava.* **2-3**:36-38 (Russ.).
- Ramsay, C.N., Ellis, P.M., and Zealley, H., 1991, Down's syndrome in the Lothian region of Scotland – 1978 to 1989, *Biomed. Pharmacother.* **45**:267-272.
- Roux, C., Horvath, C., and Dupuis, R., 1983, Effects of pre-implantation low-dose radiation on rat embryos, *Health Phys.* **45**:993-994.
- Rugh, R., and Grupp, E., 1959, X-irradiation exencephaly, *Am. J. Roentgenol. Radium Ther.Nucl. Med.* **81**:1026-1052.
- Rugh, R., 1962, Major radiobiological concepts and effects of ionizing radiations on the embryo and fetus, in Haley, T.J. and Snider, R.S., ed., *Response of the nervous system to ionizing radiation*, Academic press, New York, London, p. 3-26
- Salomaa, S., Sevan`kaev, A.V., Zhloba, A.A., Kumpusalo, E., Mäkinen, S., Lindholm, C., Kumpusalo, L., Kolmakow, S., and Nissinen, N., 1997, Unstable and stable chromosomal aberrations in lymphocytes of people exposed to Chernobyl fallout in Bryansk, Russia, *Int. J. Radiat. Biol.* **71**:51-59.

- Savchenko, V.K., 1995, The Ecology of the Chernobyl Catastrophe. Scientific outlines of an international programme of collaborative research. *Man and the Biosphere Series Vol. 17*, UNESCO Paris, p.83.
- Scheid, W., Weber, J., Petrenko, S., and Traut, H., 1993, Chromosome aberrations in human lymphocytes apparently induced by Chernobyl fallout, *Health Phys.* **64**:531-534.
- Scherb, H., and Weigelt, E., 1999a, Spatial-temporal logistic regression of the cesium contamination and the time trends in annual stillbirth proportions on a district level in Bavaria, 1980-1993, in: *Proceedings of the 14th international workshop on statistical modelling*, H. Friedl et al., ed., Technical University Graz, Graz, Austria, pp.647-650.
- Scherb, H., Weigelt, E., and Brüske-Hohlfeld, I., 1999b, European stillbirth proportions before and after the Chernobyl accident, *Int. J. Epidemiol.* **28**:932-940.
- Scherb, H., Weigelt, E., Brüske-Hohlfeld, I., 2000a, Regression analysis of time trends in perinatal mortality in Germany, *Environ. Health Persp.* **108**:159-165.
- Scherb, H., Weigelt, E., 2000b, Spatial-temporal change-point regression models for European stillbirth data, in: *30th Ann. Meeting Europ. Soc. Radiat. Biol.*, Warszawa, Poland, August 27-31, Abstracts, p.14. <http://www.tschernobylhilfe.ffb.org/dokumente/warschau.pdf>
- Scherb, H., and Weigelt, E., 2003, Congenital malformation and stillbirth in Germany and Europe before and after the Chernobyl nuclear power plant accident, *Environ. Sci. & Pollut. Res.* **10 Special (1)**:117-125.
- Scherb, H., and Weigelt, E., 2004, Cleft lip and cleft palate birth rate in Bavaria before and after the Chernobyl nuclear power plant accident, *Mund Kiefer Gesichtschir.* **8**:106-110 (in German).
- Sevan'kaev, A.V., Tsyb, A.F., Lloyd, D.C., Zhloba, A.A., Moiseenko, V.V., Skrjabin, A.M., and Klimov, V.M., 1993, 'Rogue' cells observed in children exposed to radiation from the Chernobyl accident, *Int. J. Radiat. Biol.* **63**:361-367.
- Shidlovskii, P.R., 1992, General morbidity of the population in districts of the Brest region. *Zdravoohranenie Belorussii (Minsk)* **1**:8-11 (Russ.).
- Sperling, K., Pelz, J., Wegner, R.-D., Schulzke, I., and Struck, E., 1991, Frequency of trisomy 21 in Germany before and after the Chernobyl accident, *Biomed. Pharmacother.* **45**:255-262.
- Sperling, K., Pelz, J., Wegner, R.-D., Dörries, A., Grüters, A., and Mikkelsen, M., 1994, Significant increase in trisomy 21 in Berlin nine months after the Chernobyl reactor accident: temporal correlation or causal relation? *Brit. Med. J.* **309**:158-162
- Stephan, G., and Oestreicher, U., 1989, An increased frequency of structural chromosome aberrations in persons present in the vicinity of Chernobyl during and after the reactor accident. Is this effect caused by radiation exposure? *Mutat. Res.* **223**:7-12.
- Stephan, G., and Oestreicher, U., 1993, Chromosome investigation of individuals living in areas of Southern Germany contaminated by fallout from the Chernobyl reactor accident, *Mutat. Res.* **319**:189-196.
- Strahlentelex 55, 1989, Säuglinge starben vermehrt oder wurden tot geboren, Berlin, Germany, p. 6.
- Sychik, S.I., and Stozharov, A.N., 1999, Analysis of morbidity of children irradiated in utero as a result of Chernobyl accident, *Zdravoohranenie Belorussii (Minsk)* **6**:20-22 (Russ.).
- Sychik, S.I., and Stozharov, A.N., 1999, Estimation of the influence of prenatal irradiation on functional state of critical organs and systems in children at distant terms following the Chernobyl accident. *Radiats. Biol. Radioecol.* **39**:500-504 (Russ.).
- Ulstein, M., Jensen, T.S., Irgens, L.M., Lie, R.T., and Sivertsen, E., 1990, Outcome of pregnancy in one Norwegian county 3 years prior to and 3 years subsequent to the Chernobyl accident, *Acta Obstet. Gynecol. Scand.* **6**:277-280.
- UNSCEAR: United Nations Scientific Committee on the Effects of Atomic Radiation, 1986, Genetic and Somatic Effects of Ionizing Radiation, Report to the General Assembly, New York 1986.
- UNSCEAR: United Nations Scientific Committee on the Effects of Atomic Radiation, 1988, Sources, Effects and Risks of Ionizing Radiation, Report to the General Assembly, New York 1988.
- Vershaeve, L., Domracheva, E.V., Kuznetsov, S.A., and Nechai, V.V., 1993, Chromosome aberrations in inhabitants of Byelorussia: consequence of the Chernobyl accident, *Mutat. Res.* **287**:253-259.
- Zatsepin, I.O., Verger, P., Robert-Gnansia, E., Gaguierre, B., Khemel, R.D., Lazjuk, G.I., 2004, Cluster of Down's syndrome cases registered in January 1987 in Republic of Belarus as a possible

effect of the Chernobyl accident. Int. J. Radiat. Med., Special issue 6 (1-4), Eds. Association
“Physicians of Chernobyl”, Kiev
Ziegłowski, V., and Hemprich, A., 1999, Facial cleft birth rate in former East Germany before and
after the reactor accident in Chernobyl, *Mund Kiefer Gesichtschir.* **3**:195-199 (in German).